

8_Aggregate-level-B_Snack-Tax_Results

```
-----  
  name: <unnamed>  
  log: C:\Users\ids29\Documents\Stata\Taxes_Aggregated_snack_Results.log  
log type: text  
opened on: 16 Nov 2012, 11:42:27
```

```
.  
. foreach var of varlist PINCBAD- CHL001BAD {  
  2.  
  display "----- `var' -----"  
  3.  
  signtest a_`var' = b_`var' if Snack==1  
  4.  
}.  
----- PINCBAD -----
```

Sign test

sign	observed	expected
positive	4	29
negative	54	29
zero	0	0
all	58	58

One-sided tests:

```
Ho: median of a_PINCBAD - b_PINCBAD = 0 vs.  
Ha: median of a_PINCBAD - b_PINCBAD > 0  
Pr(#positive >= 4) =  
    Binomial(n = 58, x >= 4, p = 0.5) = 1.0000  
  
Ho: median of a_PINCBAD - b_PINCBAD = 0 vs.  
Ha: median of a_PINCBAD - b_PINCBAD < 0  
Pr(#negative >= 54) =  
    Binomial(n = 58, x >= 54, p = 0.5) = 0.0000
```

Two-sided test:

```
Ho: median of a_PINCBAD - b_PINCBAD = 0 vs.  
Ha: median of a_PINCBAD - b_PINCBAD != 0  
Pr(#positive >= 54 or #negative >= 54) =  
    min(1, 2*Binomial(n = 58, x >= 54, p = 0.5)) = 0.0000
```

Sign test

sign	observed	expected
positive	0	0
negative	0	0
zero	0	0
all	0	0

One-sided tests:

```
Ho: median of a_PINCGOOD - b_PINCGOOD = 0 vs.  
Ha: median of a_PINCGOOD - b_PINCGOOD > 0  
Pr(#positive >= 0) =  
    Binomial(n = 0, x >= 0, p = 0.5) = 1.0000  
  
Ho: median of a_PINCGOOD - b_PINCGOOD = 0 vs.  
Ha: median of a_PINCGOOD - b_PINCGOOD < 0  
Pr(#negative >= 0) =  
    Binomial(n = 0, x >= 0, p = 0.5) = 1.0000
```

8_Aggregate-level-B_Snack-Tax_Results

Two-sided test:

```

Ho: median of a_PINCGOOD - b_PINCGOOD = 0 vs.
Ha: median of a_PINCGOOD - b_PINCGOOD != 0
Pr(#positive >= 0 or #negative >= 0) =
min(1, 2*Binomial(n = 0, x >= 0, p = 0.5)) = 1.0000
----- ININCBAD -----

```

Sign test

sign	observed	expected
positive	0	6
negative	12	6
zero	0	0
all	12	12

One-sided tests:

```

Ho: median of a_ININCBAD - b_ININCBAD = 0 vs.
Ha: median of a_ININCBAD - b_ININCBAD > 0
Pr(#positive >= 0) =
Binomial(n = 12, x >= 0, p = 0.5) = 1.0000

Ho: median of a_ININCBAD - b_ININCBAD = 0 vs.
Ha: median of a_ININCBAD - b_ININCBAD < 0
Pr(#negative >= 12) =
Binomial(n = 12, x >= 12, p = 0.5) = 0.0002

```

Two-sided test:

```

Ho: median of a_ININCBAD - b_ININCBAD = 0 vs.
Ha: median of a_ININCBAD - b_ININCBAD != 0
Pr(#positive >= 12 or #negative >= 12) =
min(1, 2*Binomial(n = 12, x >= 12, p = 0.5)) = 0.0005
----- ININCGOOD -----

```

Sign test

sign	observed	expected
positive	0	0
negative	0	0
zero	0	0
all	0	0

One-sided tests:

```

Ho: median of a_ININCG~D - b_ININCGOOD = 0 vs.
Ha: median of a_ININCG~D - b_ININCGOOD > 0
Pr(#positive >= 0) =
Binomial(n = 0, x >= 0, p = 0.5) = 1.0000

Ho: median of a_ININCG~D - b_ININCGOOD = 0 vs.
Ha: median of a_ININCG~D - b_ININCGOOD < 0
Pr(#negative >= 0) =
Binomial(n = 0, x >= 0, p = 0.5) = 1.0000

```

Two-sided test:

```

Ho: median of a_ININCG~D - b_ININCGOOD = 0 vs.
Ha: median of a_ININCG~D - b_ININCGOOD != 0
Pr(#positive >= 0 or #negative >= 0) =
min(1, 2*Binomial(n = 0, x >= 0, p = 0.5)) = 1.0000
----- BWINCBAD -----

```

Sign test

sign	observed	expected
positive	1	7.5

8_Aggregate-level-B_Snack-Tax_Results		
negative	14	7.5
zero	0	0
all	15	15

One-sided tests:

Ho: median of a_BWINCBAD - b_BWINCBAD = 0 vs.
 Ha: median of a_BWINCBAD - b_BWINCBAD > 0
 $\Pr(\#\text{positive} \geq 1) = \text{Binomial}(n = 15, x \geq 1, p = 0.5) = 1.0000$

Ho: median of a_BWINCBAD - b_BWINCBAD = 0 vs.
 Ha: median of a_BWINCBAD - b_BWINCBAD < 0
 $\Pr(\#\text{negative} \geq 14) = \text{Binomial}(n = 15, x \geq 14, p = 0.5) = 0.0005$

Two-sided test:

Ho: median of a_BWINCBAD - b_BWINCBAD = 0 vs.
 Ha: median of a_BWINCBAD - b_BWINCBAD != 0
 $\Pr(\#\text{positive} \geq 14 \text{ or } \#\text{negative} \geq 14) = \min(1, 2 * \text{Binomial}(n = 15, x \geq 14, p = 0.5)) = 0.0010$

Sign test

sign	observed	expected
positive	0	0
negative	0	0
zero	0	0
all	0	0

One-sided tests:

Ho: median of a_CHL001~D - b_CHL001BAD = 0 vs.
 Ha: median of a_CHL001~D - b_CHL001BAD > 0
 $\Pr(\#\text{positive} \geq 0) = \text{Binomial}(n = 0, x \geq 0, p = 0.5) = 1.0000$

Ho: median of a_CHL001~D - b_CHL001BAD = 0 vs.
 Ha: median of a_CHL001~D - b_CHL001BAD < 0
 $\Pr(\#\text{negative} \geq 0) = \text{Binomial}(n = 0, x \geq 0, p = 0.5) = 1.0000$

Two-sided test:

Ho: median of a_CHL001~D - b_CHL001BAD = 0 vs.
 Ha: median of a_CHL001~D - b_CHL001BAD != 0
 $\Pr(\#\text{positive} \geq 0 \text{ or } \#\text{negative} \geq 0) = \min(1, 2 * \text{Binomial}(n = 0, x \geq 0, p = 0.5)) = 1.0000$

```
: log close
  name: <unnamed>
  log: C:\Users\ids29\Documents\Stata\Taxes_Aggregated_Snack_Results.log
log type: text
closed on: 16 Nov 2012, 11:42:28
```